



**Manchester
Metropolitan
University**

Šimunič, Boštjan, Koren, Katja, Rittweger, Jörn, Lazzer, Stefano, Reggiani, Carlo, Rejc, Enrico, Pišot, Rado, Narici, Marco and Degens, Hans (2019) Tensiomyography detects early hallmarks of bed-rest-induced atrophy before changes in muscle architecture. *Journal of applied physiology*, 126 (4). pp. 815-822. ISSN 8750-7587

Downloaded from: <https://e-space.mmu.ac.uk/622479/>

Version: Accepted Version

Publisher: American Physiological Society

DOI: <https://doi.org/10.1152/jappphysiol.00880.2018>

Please cite the published version

<https://e-space.mmu.ac.uk>

1 **Title page**

2 **Title:** Tensiomyography detects early hallmarks of bed-rest-induced atrophy before changes in
3 muscle architecture

4

5 **Authors:**

6 Boštjan Šimunič ^{a,b}

7 Katja Koren ^b

8 Jörn Rittweger ^{c, d}

9 Stefano Lazzer ^{e, f}

10 Carlo Reggiani ^{b, g}

11 Enrico Rejc ^h

12 Rado Pišot ^b

13 Marco Narici ^{b, g}

14 Hans Degens ^{i, j, k}

15

16 **Departments and institutions:**

17 ^a University of Primorska, Titov trg 4, 6000 Koper, Slovenia

18 ^b Science and Research Centre Koper, Institute for Kinesiology Research, Garibaldijeva 1, 6000
19 Koper, Slovenia

^c Institute of Aerospace Medicine, German Aerospace Center (DLR), Linder Höhe 1, D-51147
Cologne, Germany

^d Department of Pediatrics and Adolescent Medicine, University of Cologne, Cologne,
Germany

^e Department of Medicine, University of Udine, Udine, Italy

^f School of Sport Sciences, University of Udine, Udine, Italy

^g Department of Biomedical Sciences, University of Padova, via Marzolo 3, 35131 Padova,
Italy

^h Kentucky Spinal Cord Injury Research Center, University of Louisville, Louisville, KY, USA.

ⁱ School of Healthcare Science, Manchester Metropolitan University, UK

^j Institute of Sport Science and Innovations, Lithuanian Sports University, Kaunas, Lithuania

^k University of Medicine and Pharmacy of Targu Mures, Romania

Abbreviated title:

TMG-BEDREST

Corresponding author:

Boštjan Šimunič ^{a,b}

Address: Science and Research Centre Koper, Institute for Kinesiology Research, Garibaldijeva
1, 6000 Koper, Slovenia

Email: bostjan.simunic@zrs-kp.si

41 Phon e: +38631832016

42

43 **Brief itemized list of how each author contributed to the study:**

- 44 • Study concept and design: BŠ, JR, SL, RP, MN, HD
- 45 • Acquisition of data: BŠ, SL, ER
- 46 • Analysis and interpretation of data: BŠ, KK, JR, CR, RP, MN, HD
- 47 • Drafting of the manuscript: BŠ, JR, SL, CR, ER, MN, HD
- 48 • Critical revision of the manuscript for important intellectual content: BŠ, JR, SL,
- 49 ER, RP, MN, HD

50

Abstract

In young and older people skeletal muscle mass is reduced after as little as seven days of disuse. The declines in muscle mass after such short periods are of high clinical relevance, particularly in older people who show higher atrophy rate, and a slower, or even a complete lack of muscle mass recovery after disuse. Ten men (24.3 ± 2.6 years) underwent 35 days of 6° head-down tilt bed rest followed by 30 days of recovery. During bed rest, a neutral energy balance was maintained, with three weekly passive physiotherapy sessions to minimise muscle soreness and joint stiffness. All measurements were performed in a hospital at days 1-10 (BR1-BR10), day 16 (BR16), 28 (BR28) and 35 (BR35) of bed rest, and day 1 (R+1), 3 (R+3) and 30 (R+30) after reambulation. Vastus medialis obliquus (VMO), vastus medialis longus (VML) and biceps femoris (BF) thickness (d) and pennation angle (Θ) were assessed by ultrasonography, while twitch muscle belly displacement (Dm) and contraction time (Tc) were assessed with tensiomyography. After bed rest, d and Θ decreased by 13–17% in all muscles ($P < .001$) and had recovered at R+30. Dm was increased by 42.3–84.4% ($P < .001$) at BR35 and preceded the decrease in d by 7, 5 and 3 days in VMO, VML and BF, respectively. Tc increased only in BF (32.1%; $P < .001$) and was not recovered at R+30. Tensiomyography can detect early bed-rest-induced changes in muscle with higher sensitivity before overt architectural changes and atrophy can be detected.

Key words: tensiomyography, contraction time, skeletal muscle, rehabilitation, ageing

New & Noteworthy: Detection of early atrophic processes and irreversible adaptation to disuse is of high clinical relevance. Using Tensiomyography we detected early atrophic processes before overt architectural changes and atrophy can be detected using imaging technique. Furthermore, Tensiomyography detected irreversible changes of biceps femoris contraction time.

Introduction

Hospitalization due to injury or disease can lead to a period of forced inactivity. In those conditions skeletal muscle disuse is followed by atrophy, which in turn implies loss of contractile performance and metabolic dysregulation(30). Microgravity during space flight and the experimental models of disuse have a similar impact on muscle mass and function. Studies in young adults documented that skeletal muscle mass and strength are reduced after as little as seven days of spaceflight(20, 26) or bed rest(12) and continue to decline with the length of exposure(1). Declines in muscle mass and function after such short periods are of high clinical relevance to most patients who are, on average, hospitalized for <7 days(15). The disuse-induced loss of muscle mass is particularly relevant for elderly who show higher atrophy after 14-day bed rest and a much slower recovery or even complete lack of recovery for at least 14 days afterwards (33, 36). Therefore, there is a substantial need to develop methods to detect early stages of muscle atrophy related processes.

Evidences exist that muscle atrophy is not symmetrical throughout the muscle mass. Antigravity muscles show the greatest atrophy, and distal muscles atrophy more than proximal muscles(8). In addition, muscles with different functional roles across different joints and even muscles across the same joint may respond differently to unloading(3, 8). Rehabilitation programmes and assessments after any period of disuse should thus primarily focus on postural muscles and, at the same time, not overlook the non-postural muscles(8, 49).

At the human single muscle fibre level, evidence suggests that type I fibres depict stronger atrophy in bed rest than type II muscle fibres both after bed rest(6, 7) and spaceflight(17). Furthermore, there is a slow-to-fast myosin isoform transition after bed rest(31, 45) and spaceflight(50) that would result in faster contractile properties of the muscle, which will be accentuated by an increase in maximal shortening velocity of both type I and II muscle fibres after 17-day bed rest(48) and 17-day spaceflight(47). The latter effect seems reversed after

42(25) and 84 days bed rest(45), as well as after 180 days spaceflight(17). At the whole muscle level it has been reported that the time to peak twitch isometric tension of the triceps surae muscles was increased by 13%, indicating a slowing of the musculotendinous system after 120 days of bed rest(22). However, in this latter case, this was attributable to reduced tendon stiffness and increased muscle-tendon passive elasticity(23, 35), and thus not due to alterations in muscle contractile properties.

While ultrasound provides a reliable and non-invasive tool to follow structural changes of skeletal muscle during disuse, functional assessment of e.g. twitch torque requires specialised equipment and may not always be possible in bed ridden patient(21, 34, 38, 39, 41). To overcome this problem, relatively simple and low cost mechanomyographic methods were developed, where for instance Tensiomyography (TMG) allows for non-invasive and reliable(38, 44) estimation of contraction time (T_c), selectively in superficial muscle heads. This method can estimate the percentage of type I myosin heavy chain at least in the vastus lateralis (VL) muscle(39), and possibly also in other muscles. There is a clear distinction between results obtained from twitch torque and TMG. For example, the T_c is 42.7% shorter when estimated from TMG than from twitch torque(21). This indirectly confirms that TMG gives better insights to the muscle contractility as it is less affected by the surrounding tissues(16, 21).

Using TMG, it was found that after 35 days of bed rest there was no change in T_c of the vastus medialis, but an increased T_c in gastrocnemius medialis muscle(34). The authors did, however, report that the TMG amplitude (D_m) was increased in both muscles, and that for gastrocnemius medialis the change in D_m was negatively correlated to the change in thickness ($r=-.70$). The D_m increase in both muscles in the abovementioned study may indicate a lower muscle resting tension and, possibly, decreased visco-elasticity(16).

While TMG detects changes after a prolonged disuse period(34), nothing is known about the possibility to adopt this method to follow initial and early changes in the adaptive response of muscle to disuse, before overt measurable atrophy. Therefore, the aim of our study was to assess 1) the time course of changes in muscle architecture and TMG parameters during 35 days bed rest and the following 30 days supervised recovery in young men, and 2) whether TMG is able to detect early changes that occur just after a few days of disuse.

Methods

Participants

Ten healthy men (age: 24.3 ± 2.6 years, Table 1) with no history of neuromuscular or cardiovascular disorders participated in our study. The study was approved by the Slovenian National Medical Ethics Committee (approval number 72/06/08). All participants were fully informed about the study procedures and the possible health risks of study participation. Routine medical and laboratory analyses were performed to exclude participants with chronic diseases. None of the subjects regularly took any medication. From all participants written informed consent was obtained prior the study. All procedures were in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki and its amendments.

<< Insert Table 1 >>

Experimental design

The bed rest study was conducted in the Orthopaedic hospital of Valdoltra under medical supervision. Participants arrived a week before the bed rest and were asked to visit the

laboratory on several occasions to become familiar with testing procedures. All baseline data were collected (BDC) 1 day before the start of bed rest. After BDC, participants went through 35 days 6° head-down tilt bed rest followed by 30 days of supervised recovery. Subsequent measurements were performed at days 1-10 (BR1-BR10), day 16 (BR16), 28 (BR28) and 35 (BR35) of bed rest, and day 1 (R+1), 3 (R+3) and 30 (R+30) after completion of bed rest. During recovery, a fitness professional was available and all participants received written recovery instructions. Recovery consisted of 12 sessions (3 sessions/week). Each session lasted about 60 minutes and consisted of a 10-min warm-up, 5 min active stretching, followed by 20 min strength and balance exercises and 20 min aerobic exercises and a 5-min cool-down.

During bed rest, the participants received three weekly passive physiotherapy sessions to minimise muscle soreness and joint stiffness. Each participant received a weight-maintaining diet with an energy content of 1.4 and 1.2 times his resting energy expenditure, calculated using the FAO/WHO equations(29), for the pre-bed rest and bed rest period, respectively(5). The diet contained 60% of energy as carbohydrate, 25% as fat and 15% as protein. Six meals were administered daily: 3 main meals (breakfast, lunch and dinner) and 3 snacks. Subjects were required to consume all food served.

Measurements

Ultrasonography

Muscle architecture was determined at rest with B-mode ultrasonography (Mylab 25, 13-4 MHz, linear array transducer probe LA523, Esaote Biomedica, Geneva, Italy). Biceps femoris (BF) scans were taken with the participant prone and with a knee angle set at 5° flexion with foam pads. The BF measuring site was halfway between the ischial tuberosity and the posterior knee joint fold, along the line of the BF long head. Vastus medialis obliquus (VMO) scans were

obtained supine at a knee angle set at 30° flexion with foam pads. The VMO measuring site was at the midpoint of the line from the patella to the VMO innervation point. The vastus medialis longus (VML) scans were obtained supine at 30° knee flexion at the midpoint of the line from the patella to the VML innervation point. The VMO and VML innervation points were detected using monophasic tetanic stimulation (impulse width 0.1 ms; frequency 10 Hz). To ensure that all subsequent ultrasound measurements were taken at the same anatomical location, the ultrasound probe was positioned in the midsagittal plane, orthogonal to the mediolateral axis, and its positioning was marked on acetate paper using moles and small angiomas as reference points.

For each muscle, three scans were obtained. Thickness (d in mm) and pennation angle (Θ in °) were measured using Matlab (Matlab, The MathWorks Inc., USA). In each scan, the fascicular path was determined as the interspaces between echoes coming from the perimysial tissue surrounding the fascicle. Muscle thickness was defined as the shortest distance between the deep and superficial aponeuroses. Pennation angle was defined as the angle between the fascicle pathway and the deep aponeurosis of the muscle. The average values for each architecture parameter of three scans was used for further analysis

Tensiomyography

Tensiomyography (TMG) was assessed in the same muscles at the same body positions and at the same measurement sites as ultrasound scans. TMG measurements were performed during electrically-evoked maximal isometric contractions. A single 1-ms maximal monophasic electrical impulse was used to elicit a twitch contraction that caused the muscle belly to oscillate. These oscillations were recorded using a sensitive digital displacement sensor (TMG-BMC Ltd., Ljubljana, Slovenia) that was placed on the surface of the skin at the measuring site

of the muscle of interest. Initially, the stimulation amplitude was set just above the threshold and then gradually increased until the amplitude of the radial twitch displacement (D_m in mm) increased no further. Electrical pulses ranged between 85 and 110 milliamperes at constant 30 volts. From two maximal twitch responses, also contraction time (T_c in ms) was calculated (Figure 1) as the time for the amplitude to increase from 10% to 90% of D_m (Figure 1)(39, 42). Furthermore, the velocity of radial displacement (V_r) was calculated by dividing $.8 \cdot D_m$ with T_c (37).

<< Insert Figure 1 here >>

Statistics

SPSS (IBM Ltd., USA) software was used for all statistical analyses. All data in text and tables are presented as mean \pm standard deviation, while in figures standard errors were used. Visual inspection and the Shapiro-Wilk test indicated that all data were normally distributed. Sphericity (homogeneity of covariance) was verified by the Mauchly's test. When the assumption of sphericity was not met, the significance of the F-ratios was adjusted according to the Greenhouse-Geisser procedure. Main effects were studied with a General Linear Model repeated-measures ANOVA with time (BDC, BR_i, R+j; where $i = 1-10, 16, 28, 35$ and $j = 1, 3, 30$) and muscle (VMO, VML, BF) as within factors. If a significant time x muscle interaction was found, the analysis was repeated with relative data representing percent change from BDC, to exclude any bias related to e.g. a difference in muscle thickness at BDC between muscles. Where significant time, muscle and interaction time x muscle effects were found, post-hoc analysis with Bonferroni corrections was used to locate the differences in time ($p' = p/16$; where 16 is the number of comparisons to the BDC value) for each muscle. Pearson regression analysis

was used to correlate changes during bed rest ($\Delta(\text{BDC}-\text{BR35})$) in Tc and Dm to changes in muscle architecture. Statistical significance was accepted at $p \leq .05$. The effect size for dependent variables was given as partial eta-squared (η^2).

Results

The variations in muscle structure as determined by ultrasonography and of muscle contractile function as measured with TMG are reported in Figure 2. Skeletal muscle thickness changed during the study ($P < .001$; $\eta^2 = .865$; Figure 2A). Specifically, thickness declined progressively by 4.5% at BR7 ($P = .048$) to 15.2% at BR35 ($P < .001$), and recovered to BDC thickness at R+30 ($P = .22$). The absence of a time x muscle interaction ($P = .50$), indicates that the % changes in muscle thickness during bed rest and recovery did not differ significantly between muscles.

The time x muscle interaction ($P < .001$; $\eta^2 = .938$) for Θ indicates that the changes in Θ over time differed between the three muscles. While the time course was qualitatively similar for the three muscles ($P < .001$; $\eta^2 = .592$; Figure 2B), post-hoc analysis revealed that in the VMO Θ was first significantly decreased at BR6 (13.6%; $P = .033$), while in VML and BF it was already decreased at BR2 (5.5%; $P = .037$) and BR3 (7.4%; $P = .019$), respectively, interestingly at smaller decrease due to lower variance. In VMO and VML Θ had recovered to BDC at R+30 ($P > .05$) while in BF it was already recovered at R+3 ($P = .32$).

Two parameters characterize the TMG signal, the Dm and Tc, as well as the ratio between them, the V_r . The muscle x time interaction for Dm ($P < .001$; $\eta^2 = .186$) indicates that the changes in Dm during the study ($P < .001$; $\eta^2 = .782$; differed between the three muscles (Figure 2C). While the time course was qualitatively similar for the muscles, the magnitude of the rise in Dm was larger in the VML (84.4%) and BF (75.6%) than in the VMO (42.3%) at BR35 ($P = .013$;

$\eta^2=.381$). Dm increased already after BR1, BR4 and BR6 in VMO, VML and BF, respectively, and had returned to BDC at R+3 ($P=.050$).

The muscle x time interaction for Tc ($P<.001$; $\eta^2=.255$) indicates that the changes in Tc during the study ($P<.001$; $\eta^2=.397$; Figure 2D) differed between the three muscles. Post-hoc analysis revealed that Tc of the VMO did not change significantly during bed rest and recovery ($P=.35$), while the Tc of the VML ($P<.001$; $\eta^2=.300$) and BF ($P<.001$; $\eta^2=.393$) did change. We were unable to locate the difference with post-hoc tests in the VML. In the BF we found an increased Tc at BR7 (23.6% $P=.043$), being highest at R+1 (39.3%; $P=.013$). BF Tc did not return to the BDC value even at R+30 (26.4%; $P=.041$).

The muscle x time interaction for V_r ($P<.001$; $\eta^2=.283$) indicates that the changes in V_r during the study ($P<.001$; $\eta^2=.733$; Figure 2E) differed between the three muscles. We found differences in V_r at BDC ($P=.017$), where V_r was slowest in BF in comparison to VM muscles ($P=.014$). Furthermore, post-hoc analysis revealed that V_r of the VMO, VML and BF increased during bed rest for 40.7% ($P<.001$; $\eta^2=.609$) after BR9, for 74.6% ($P<.001$; $\eta^2=.679$) after BR6 and for 36.1% ($P<.001$; $\eta^2=.418$) after BR16, respectively. In all muscles V_r returned to BDC at R+1.

<< Insert Figure 2 here >>

The contractile parameters measured with TMG and the structural parameters measured with ultrasonography revealed correlations (Figure 3). Changes in muscle thickness and Dm between BDC and BR35 were negatively correlated. This negative correlation was significant in the BF ($P=.001$), but not in the VMO ($P=.09$) and VML ($P=.06$). There was also a positive correlation between Dm and Θ in VMO ($P=.008$) and VML ($P=.050$).

266

267

<< Insert Figure 3 here >>

268

269 Changes in Tc did not correlate significantly with changes in any of the architectural parameters
270 (data not shown).

271

272 Discussion

273 Thirty-five days of 6° head-down bed rest induced a similar degree of atrophy (reduction in
274 thickness) across all three muscles that had recovered 30 days after completion of bed rest. The
275 atrophy was accompanied by a reduction in Θ that returned to baseline levels as soon as 3 days
276 after cessation of bed rest. While the degree of atrophy became significant only after 7 days of
277 bed rest, the increase in Dm was significant as soon as 1, 4 and 6 days after initiation of bed
278 rest in the VMO, VML and BF, respectively. This suggests that Dm determined by TMG can
279 be used to non-invasively and easily detect early hallmarks of the atrophy process, before overt
280 atrophy was measurable by ultrasound.

281 After 35 days bed rest the muscle thickness was decreased by 16-23%, which is similar to the
282 amount of atrophy seen in other studies(2, 4, 8, 28). In contrast to other studies(2, 4, 8, 28), we
283 did not observe differences in the relative degree of atrophy between muscles. The discrepancy
284 between these studies and ours may well be related to the range of muscles studied, where we
285 assessed the bed-rest-induced changes only in the thigh, where others have compared the thigh
286 muscles with muscles in the lower leg that atrophied more. It is likely that this difference in
287 bed-rest-induced decreases in muscle mass between muscles is related to a larger reduction in

recruitment of lower leg than thigh muscles during bed rest. As expected, the atrophy was accompanied by a decline in Θ in all muscles as was previously also demonstrated(8).

Similar to a previous study we found that in all muscles Dm was increased by 35 days of bed rest though the increase in the present study was more pronounced than in that study using horizontal bed rest(34). This suggests that the fluid shift, away from the legs towards the head somehow affects the atrophy-induced increase in Dm. The fluid shift may also contribute to the observation that Dm was already elevated after as little as 24 hours of bed rest, before any overt architectural changes and muscle atrophy had taken place. In addition, the magnitude of Dm increase was between 42 and 84% after 35 days head-down tilt bed-rest and exceeded the atrophy that ranged between 16 and 23%. Another indicator that the fluid shift may play an important role in the increase in Dm with bed rest, is the almost instantaneous return of Dm after cessation of bed rest (at R+3), again before any significant architectural and muscle mass recovery had taken place (at R+30, except Θ in BF at R+3). How the fluid shift affects these changes is a matter of further research, but one might speculate that Dm may also be applicable to assess the hydration status of the muscle.

It is possible that fluid shifts out of the muscle may increase Dm by decreasing the viscoelasticity of the muscle-tendon tissue and decrease in muscle tone, resulting a larger bulging of the muscle in response to an identical electrical stimulus. The fluid shift from extremities to the chest can amount to a 4.4% decrease in extracellular fluid content that is particularly attributable to a loss of interstitial volume by 3% in parallel with a 12.3% reduction in plasma volume in just 4 days(19). After the 4th day of bed rest plasma volume continues to decrease, but at a much slower rate(19). Later also intracellular fluid loss can occur that then parallels muscle atrophy(19).

Also dry immersion induces an increase in Dm and decrease muscle-tendon viscoelasticity(10, 24), that is at least partly attributable to a similar fluid shift away from the muscles. A decrease

in muscle tone, which occurs as early as after 1 day of dry immersion, may further contribute to the increased Dm after 3 days of dry immersion(10) and after 20 days of bed rest(24). Such changes have indeed been observed to translate into higher transversal muscle oscillations during voluntary and electrically-evoked contractions(27).

Recent data show that merely a few days (e.g. 5-7 days) of disuse substantially reduce skeletal muscle mass(11, 13), with a slower recovery rate in seniors than in young adults(33, 43). As a consequence, it has been suggested that the accumulation of such short (<10 days), successive periods of bed rest or immobilization during short-term illness or hospitalisation may contribute to the loss of muscle mass and metabolic decline observed throughout life(14, 46). Given this slow recovery in the older person, and being more prone to hospitalisation, it is important to minimise, or even prevent, any atrophy. Identification of early functional and structural markers of muscle deconditioning may help in designing adequate interventions to slow such atrophy even before it becomes overt, and assess the success of an intervention to prevent atrophy(10). Our data show that Dm may be such a functional marker, a parameter that can be determined with high reproducibility(38, 44).

Bed rest did not induce a significant change in the Tc in the VMO, but did induce an increase in the Tc in the VML and BF muscles. That observed increase was much more pronounced for BF, where Tc also did not recover until 30 days after bed rest. Previously we found a positive correlation between Tc and the MHC-I proportion in VL(39), and given that disuse is often associated with a slow MHC-I to fast MHC-IIx transition, the correlation may not apply to disused muscles, where for instance a decreased visco-elasticity may have a larger, and opposite to, effect than the myosin heavy chain transition. However, the velocity of radial displacement (Vr) increased in all muscles resulting from increased Dm and: (i) unchanged Tc in VMO, (ii) slightly increased Tc in VML; and (iii) substantially increased Tc in BF. Although Vr should not be paralleled to the contractile velocity of the whole muscle it is evident that Vr is sensitive

to muscle disuse as well as to assess peripheral fatigue after training(32) or peripheral arterial disease(18). However, further research is needed for the interpretation of Vr changes. Whatever the explanation, the data are analogous to the lower TMG-derived Tc in children and adults who participated regularly in sports(40, 42), or high-speed plyometric exercise(51). Indeed, when compared to previously published data, the magnitude of the increase in Tc after 35 days bed rest was comparable to or even more pronounced than that of sedentary childhood/adolescence or sedentary ageing (Table 2).

The increase in Tc in the BF following bed rest may have significant implications as it has been observed that a lower Tc correlated to higher vertical jump(51). The increase in Tc following bed rest in the BF, that was found also in seniors(42) may thus have significant clinical implications for the quality of life after hospitalisation. Therefore Tc of the BF is a parameter, like Dm, of special interest in assessing the efficacy of therapeutical interventions of people going through any kind of disuse, especially in the older population(33, 43).

<< Insert Table 2 >>

Conclusions/Relevance

In conclusion, our study showed that TMG can be used to detect early bed-rest-induced muscle dysfunction, before overt atrophy and atrophy-associated architectural changes can be detected with ultrasound. It remains to be seen whether such early changes are a result of the fluid shift away from muscle during head-down bed rest and/or is a reflection of structural bed-rest induced changes. Future studies in horizontal bed rest or unilateral limb suspension may shed light on the role of fluid shifts in TMG parameters. If no such changes are observed in such a model it is probably worthwhile to assess whether TMG can be used as clinical diagnostic tool

for atrophy and/or to assess the hydration status, something particularly important in older people and chronically ill patients where dehydration is related to sarcopenia and muscle weakness(9).

Acknowledgements

We are thankful to Agenzia spaziale Italiana -ASI for financing this study (Grant No. ASI N.I /045/08/0). We are also thankful to study participants, Orthopaedic hospital Valdoltra staff for their willingness to participate in the study.

Conflict of interest statement: There are no conflicts of interest.

373 **Table 1:** Anthropometric data of participants.

	BDC	BR35	R+30	P (η^2)
N	10			
Body height / m	1.78±6.5	1.78±6.5 ¹	1.78±6.6	.92
Body mass / kg	75.3±9.3	72.2±8.7 [‡]	74.8±8.2	<.001 (.709)
Fat mass / kg	15.8±3.6	15.7±3.2	14.4±2.6 [†]	.003 (.470)
Body mass index / kg/m ²	23.7±1.9	22.7±1.7 [‡]	23.6±1.7	<.001 (.700)

374 *Values are means ± SD; BDC: Before bed rest; BR35: 35 days bed rest; R+30: after 30 days*
 375 *recovery; ¹ body height was measured 12 hours after reambulation; * P<.05; [†] P<.01; [‡] P<.001*
 376 *significantly different from BDC.*

377

378 **Table 2:** Biceps femoris contraction time of men: data from different populations/studies.

Population	N	Contraction time / ms	Reference
Children and adolescents			(40)
10 years – pooled	53	30.8 ± 5.0	
14 years – pooled	53	31.9 ± 6.3	
14 years – sedentary group	17	35.3 ± 9.1	
14 years – athletes	29	30.7 ± 6.1	
Adults (24 years)	10		This study
Before bed rest		28.3 ± 7.4	
After 35-day bed rest		36.1 ± 6.1	
After 30-day re-training		34.7 ± 6.9	
Adults, students (22 years)	20		(51)
Before plyometrics		30.6 ± 7.7	
After 8 weeks of plyometrics		24.7 ± 5.9	
Adults and seniors			(42)
35-49 years – power master athletes	32	26.6 ± 7.0	
35-49 years – sedentary group	31	33.5 ± 7.0	
35-49 years – endurance master athletes	20	41.0 ± 8.5	
50-64 years – power master athletes	33	34.3 ± 8.9	
50-64 years – sedentary group	45	41.5 ± 11.4	
50-64 years – endurance master athletes	25	40.1 ± 6.5	
65+ years – power master athletes	35	38.9 ± 9.0	
65+ years – sedentary group	57	44.3 ± 9.2	
65+ years – endurance master athletes	31	53.4 ± 10.5	

379

380 *Values are means \pm SD*

381

Literature

1. **Adams GR, Caiozzo VJ, Baldwin KM.** Skeletal muscle unweighting: spaceflight and ground-based models. *J Appl Physiol* 95: 2185–2201, 2003.
2. **Alkner BA, Tesch PA.** Efficacy of a gravity-independent resistance exercise device as a countermeasure to muscle atrophy during 29-day bed rest. *Acta Physiol Scand* 181: 345–357, 2004.
3. **Belavý DL, Miokovic T, Armbrrecht G, Richardson CA, Rittweger J, Felsenberg D.** Differential atrophy of the lower-limb musculature during prolonged bed-rest. *Eur J Appl Physiol* 107: 489–499, 2009.
4. **Belavý DL, Ohshima H, Rittweger J, Felsenberg D.** High-intensity flywheel exercise and recovery of atrophy after 90 days bed--rest-. *BMJ Open Sport Exerc Med* 3: e000196, 2017.
5. **Biolo G, Agostini F, Šimunič B, Sturma M, Torelli L, Preiser JC, Deby-Dupont G, Magni P, Strollo F, Di Prampero P, Guarnieri G, Mekjavič IB, Pišot R, Narici M V.** Positive energy balance is associated with accelerated muscle atrophy and increased erythrocyte glutathione turnover during 5 wk of bed rest. *Am J Clin Nutr* 88: 950–958, 2008.
6. **Blottner D, Bosutti A, Degens H, Schiffel G, Gutschmann M, Buehlmeier J, Rittweger J, Ganse B, Heer M, Salanova M.** Whey protein plus bicarbonate supplement has little effects on structural atrophy and proteolysis marker immunopatterns in skeletal muscle disuse during 21 days of bed rest. [Online]. *J Musculoskelet Neuronal Interact* 14: 432–44, 2014. <http://www.ncbi.nlm.nih.gov/pubmed/25524969>.
7. **Blottner D, Salanova M, Püttmann B, Schiffel G, Felsenberg D, Buehring B,**

- Rittweger J.** Human skeletal muscle structure and function preserved by vibration muscle exercise following 55 days of bed rest. *Eur J Appl Physiol* 97: 261–271, 2006.
8. **de Boer MD, Seynnes OR, di Prampero PE, Pišot R, Mekjavić IB, Biolo G, Narici M V.** Effect of 5 weeks horizontal bed rest on human muscle thickness and architecture of weight bearing and non-weight bearing muscles. *Eur J Appl Physiol* 104: 401–407, 2008.
9. **Degens H, Wüst RCI.** Water: The fountain of strength. *Acta Physiol.* (June 21, 2018). doi: 10.1111/apha.13153.
10. **Demangel R, Treffel L, Py G, Briocche T, Pagano AF, Bareille MP, Beck A, Pessemesse L, Candau R, Gharib C, Chopard A, Millet C.** Early structural and functional signature of 3-day human skeletal muscle disuse using the dry immersion model. *J Physiol* 595: 4301–4315, 2017.
11. **Dirks ML, Wall BT, Snijders T, Ottenbros CLP, Verdijk LB, Van Loon LJC.** Neuromuscular electrical stimulation prevents muscle disuse atrophy during leg immobilization in humans. *Acta Physiol* 210: 628–641, 2014.
12. **Dirks ML, Wall BT, Van De Valk B, Holloway TM, Holloway GP, Chabowski A, Goossens GH, Van Loon LJ.** One week of bed rest leads to substantial muscle atrophy and induces whole-body insulin resistance in the absence of skeletal muscle lipid accumulation. *Diabetes* 65: 2862–2875, 2016.
13. **Dirks ML, Wall BT, van de Valk B, Holloway TM, Holloway GP, Chabowski A, Goossens GH, van Loon LJC.** One Week of Bed Rest Leads to Substantial Muscle Atrophy and Induces Whole-Body Insulin Resistance in the Absence of Skeletal Muscle Lipid Accumulation. *Diabetes* 65: 2862–2875, 2016.

- 428 14. **English KL, Paddon-Jones D.** Protecting muscle mass and function in older adults
429 during bed rest. *Curr Opin Clin Nutr Metab Care* 13: 34–9, 2010.
- 430 15. **Eurostat.** Hospital discharges and length of stay statistics [Online]. *Stat. Explain.*
431 *Hosp. discharges length Stay Stat.*: 2017. [http://ec.europa.eu/eurostat/statistics-](http://ec.europa.eu/eurostat/statistics-explained/index.php/Hospital_discharges_and_length_of_stay_statistics)
432 [explained/index.php/Hospital_discharges_and_length_of_stay_statistics](http://ec.europa.eu/eurostat/statistics-explained/index.php/Hospital_discharges_and_length_of_stay_statistics) [4 May 2018].
- 433 16. **Evetovich TK, Housh TJ, Stout JR, Johnson GO, Smith DB, Ebersole KT.**
434 Mechanomyographic responses to concentric isokinetic muscle contractions. *Eur J*
435 *Appl Physiol Occup Physiol* 75: 166–169, 1997.
- 436 17. **Fitts RH, Trappe SW, Costill DL, Gallagher PM, Creer AC, Colloton PA, Peters**
437 **JR, Romatowski JG, Bain JL, Riley DA.** Prolonged space flight-induced alterations
438 in the structure and function of human skeletal muscle fibres. *J Physiol* 588: 3567–
439 3592, 2010.
- 440 18. **Gasparini M, Sabovic M, Gregoric ID, Simunic B, Pisot R.** Increased fatigability of
441 the gastrocnemius medialis muscle in individuals with intermittent claudication. *Eur. J.*
442 *Vasc. Endovasc. Surg.* (2012). doi: 10.1016/j.ejvs.2012.04.024.
- 443 19. **Greenleaf JE, Stinnett HO, Davis GL, Kollias J, Bernauer EM.** Fluid and
444 electrolyte shifts in women during +Gz acceleration after 15 days' bed rest. *J Appl*
445 *Physiol* 42: 67–73, 1977.
- 446 20. **Grigoryeva, LS; Kozlovskaya I.** Effect of weightlessness and hypokinesia on velocity
447 and strength properties of human muscles. *Kosm Biol I Aviakosmicheskaya Meditsina*
448 21: 27–30, 1987.
- 449 21. **Koren K, Šimunič B, Rejc E, Lazzer S, Pišot R.** Differences between skeletal muscle
450 contractile parameters estimated from transversal tensiomyographic and longitudinal

torque twitch response. *Kinesiology* 47: 19–26, 2015.

22. **Koryak Y.** Contractile properties of the human triceps surae muscle during simulated weightlessness. *Eur J Appl Physiol Occup Physiol* 70: 344–350, 1995.

23. **Kubo K, Akima H, Ushiyama J, Tabata I, Fukuoka H, Kanehisa H, Fukunaga T.** Effects of resistance training during bed rest on the viscoelastic properties of tendon structures in the lower limb. *Scand J Med Sci Sport* 14: 296–302, 2004.

24. **Kubo K, Akima H, Ushiyama J, Tabata I, Fukuoka H, Kanehisa H, Fukunaga T.** Effects of 20 days of bed rest on the viscoelastic properties of tendon structures in lower limb muscles. *Br J Sports Med* 38: 324–330, 2004.

25. **Larsson L, Li X, Berg HE, Frontera WR.** Effects of removal of weight bearing function on contractility and myosin isoform composition in single human skeletal muscle cells. *Pflugers Arch Eur J Physiol* 432: 320–328, 1996.

26. **LeBlanc A, Rowe R, Schneider V, Evans H, Hedrick T.** Regional muscle loss after short duration spaceflight. *Aviat Sp Environ Med* 66: 1151–1154, 1995.

27. **Longo S, Cè E, Rampichini S, Devoto M, Limonta E, Esposito F.** Mechanomyogram amplitude correlates with human gastrocnemius medialis muscle and tendon stiffness both before and after acute passive stretching. *Exp Physiol* 99: 1359–1369, 2014.

28. **Miokovic T, Armbrecht G, Felsenberg D, Belavý DL.** Differential atrophy of the postero-lateral hip musculature during prolonged bedrest and the influence of exercise countermeasures. *J Appl Physiol* 110: 926–934, 2011.

29. **Müller MJ, Bosy-Westphal A, Klaus S, Kreymann G, Lührmann PM, Neuhäuser-Berthold M, Noack R, Pirke KM, Platte P, Selberg O, Steiniger J.** World Health

Organization equations have shortcomings for predicting resting energy expenditure in persons from a modern, affluent population: generation of a new reference standard from a retrospective analysis of a German database of resting energy expenditure. *Am J Clin Nutr* 80: 1379–1390, 2004.

30. **Narici M V., De Boer MD.** Disuse of the musculo-skeletal system in space and on earth. *Eur J Appl Physiol* 111: 403–420, 2011.

31. **Ohira Y, Yoshinaga T, Ohara M, Nonaka I, Yoshioka T, Yamashita-Goto K, Shenkman BS, Kozlovskaya IB, Roy RR, Edgerton VR.** Myonuclear domain and myosin phenotype in human soleus after bed rest with or without loading. *J Appl Physiol* 87: 1776–1785, 1999.

32. **De Paula Simola R, Harms N, Raeder C, Kellmann M, Meyer T, Pfeiffer M, Ferrauti A.** Assessment of neuromuscular function after different strength training protocols using tensiomyography. *J. Strength Cond. Res.* (2015). doi: 10.1519/JSC.0000000000000768.

33. **Pišot R, Marušič U, Biolo G, Mazzucco S, Lazzer S, Grassi B, Reggiani C, Toniolo L, di Prampero PE, Passaro A, Narici M V., Mohammed S, Rittweger J, Gasparini M, Gabrijelčič Blenkuš M, Šimunič B.** Greater loss in muscle mass and function but smaller metabolic alterations in older compared with younger men following 2 wk of bed rest and recovery. *J Appl Physiol* 120: 922–929, 2016.

34. **Pišot R, Narici M V., Šimunič B, De Boer M, Seynnes O, Jurdana M, Biolo G, Mekjavič IB.** Whole muscle contractile parameters and thickness loss during 35-day bed rest. *Eur J Appl Physiol* 104: 409–414, 2008.

35. **Reeves ND.** Influence of 90-day simulated microgravity on human tendon mechanical properties and the effect of resistive countermeasures. *J Appl Physiol* 98: 2278–2286,

- 498 2005.
- 499 36. **Rejc E, Floreani M, Taboga P, Botter A, Toniolo L, Cancellara L, Narici M V.,**
500 **Šimunič B, Pišot R, Biolo G, Passaro A, Rittweger J, Reggiani C, Lazzar S.** Loss of
501 maximal explosive power of lower limbs after 2 weeks of disuse and incomplete
502 recovery after retraining in older adults. *J Physiol* 596: 647–665, 2018.
- 503 37. **Rodríguez-Ruiz D, Diez-Vega I, Rodríguez-Matoso D, Fernandez-Del-Valle M,**
504 **Sagastume R, Molina JJ.** Analysis of the response speed of musculature of the knee
505 in professional male and female volleyball players. *Biomed Res. Int.* (2014). doi:
506 10.1155/2014/239708.
- 507 38. **Šimunič B.** Between-day reliability of a method for non-invasive estimation of muscle
508 composition. *J Electromyogr Kinesiol* 22: 527–530, 2012.
- 509 39. **Šimunič B, Degens H, Rittweger J.** Noninvasive Estimation of Myosin Heavy Chain
510 Composition in Human Skeletal Muscle. *Med Sci Sport Exerc Sport Exerc* d: 27–30,
511 2011.
- 512 40. **Šimunič B, Degens H, Završnik J, Koren K, Volmut T, Pišot R.** Tensiomyographic
513 Assessment of Muscle Contractile Properties in 9- to 14-Year Old Children. *Int J*
514 *Sports Med* 38: 659–665, 2017.
- 515 41. **Šimunič B, Križaj D, Narici M V., Pišot R.** Twitch parameters in transversal and
516 longitudinal biceps brachii response. *Ann Kinesiol* 1: 61–80, 2010.
- 517 42. **Šimunic B, Pišot R, Rittweger J, Degens H.** Age-related Slowing of Contractile
518 Properties Differs between Power-, Endurance- and non-athletes; a Tensiomyographic
519 Assessment. *J Gerontol A Biol Sci Med Sci* 1, 2018.
- 520 43. **Suetta C, Frandsen U, Mackey AL, Jensen L, Hvid LG, Bayer ML, Petersson SJ,**

- 521 **Schröder HD, Andersen JL, Aagaard P, Schjerling P, Kjaer M.** Ageing is
 522 associated with diminished muscle re-growth and myogenic precursor cell expansion
 523 early after immobility-induced atrophy in human skeletal muscle. *J Physiol* 591: 3789–
 524 3804, 2013.
- 525 44. **Tous-Fajardo J, Moras G, Rodríguez-Jiménez S, Usach R, Doutres DM,**
 526 **Maffiuletti NA.** Inter-rater reliability of muscle contractile property measurements
 527 using non-invasive tensiomyography. *J Electromyogr Kinesiol* 20: 761–766, 2010.
- 528 45. **Trappe S, Trappe T, Gallagher P, Harber M, Alkner B, Tesch P.** Human single
 529 muscle fibre function with 84 day bed-rest and resistance exercise. *J Physiol* 557: 501–
 530 513, 2004.
- 531 46. **Wall BT, Dirks ML, van Loon LJC.** Skeletal muscle atrophy during short-term
 532 disuse: implications for age-related sarcopenia. *Ageing Res Rev* 12: 898–906, 2013.
- 533 47. **Widrick JJ, Knuth ST, Norenberg KM, Romatowski JG, Bain JL, Riley DA,**
 534 **Karhanek M, Trappe SW, Trappe TA, Costill DL, Fitts RH.** Effect of a 17 day
 535 spaceflight on contractile properties of human soleus muscle fibres. *J Physiol* : 915–30,
 536 1999.
- 537 48. **Widrick JJ, Norenberg KM, Romatowski JG, Blaser C a, Karhanek M, Sherwood**
 538 **J, Trappe SW, Trappe T a, Costill DL, Fitts RH.** Force-velocity-power and force-
 539 pCa relationships of human soleus fibers after 17 days of bed rest. *J Appl Physiol* 85:
 540 1949–1956, 1998.
- 541 49. **Widrick JJ, Trappe SW, Romatowski JG, Riley D a, Costill DL, Fitts RH.**
 542 Unilateral lower limb suspension does not mimic bed rest or spaceflight effects on
 543 human muscle fiber function. *J Appl Physiol* 93: 354–60, 2002.

- 544 50. **Zhou MY, Klitgaard H, Saltin B, Roy RR, Edgerton VR, Gollnick PD.** Myosin
545 heavy chain isoforms of human muscle after short-term spaceflight. *J Appl Physiol* 78:
546 1740–1744, 1995.
- 547 51. **Zubac D, Šimunič B.** Skeletal Muscle Contraction Time and Tone Decrease After 8
548 Weeks of Plyometric Training. *J strength Cond Res* 31: 1610–1619, 2017.
- 549

550 **Figure 1:** Typical tensiomyographic response of the vastus medialis obliquus (left) and biceps
551 femoris (right) at baseline (solid line) and after 35 days of bed rest (broken line).

552 Tc: contraction time defined as the time from 10% to 90% of the maximal displacement
553 amplitude (Dm).

554

555 **Figure 2:** Changes in A) thickness (d), B) pennation angle (Θ), C) tensiomyographic
556 displacement (Dm), D) contraction time (Tc) and E) velocity of radial displacement (Vr) during
557 35 days bed rest and 30 days recovery in the Vastus medialis oblique (VMO), vastus medialis
558 longus (VML) and biceps femoris (BF).

559

560 *Values are means \pm SE*

561

562 **Figure 3:** Pearson correlations between changes in tensiomyographic displacement (Dm) after
563 35 day bed rest and thickness (A-C) and pennation angle (Θ ; D-F) in the vastus medialis
564 obliquus (VMO), vastus medialis longus (VML) and biceps femoris (BF).